

Ø Question # 1

i) From the given table we have

$$0+0=0 \quad \text{and} \quad 0+1=1$$

This show that 0 is the identity element.

\oplus	0	1
0	1	1
1	1	0

ii) Since $1+1=0$ (identity element) so the inverse of 1 is 1.

iii) It is clear from table that element of the given set satisfy closure law, associative law, identity law and inverse law thus given set is group under \oplus .

Also it satisfies commutative law so it is an abelian group.

Ø Question # 2

Suppose $G = \{0,1,2,3\}$

i) The given table show that each element of the table is a member of G thus closure law holds.

ii) \oplus is associative in G .

iii) Table show that 0 is identity element w.r.t. \oplus .

iv) Since $0+0=0, 1+3=0, 2+2=0, 3+1=0$

$$\Rightarrow 0^{-1}=0, 1^{-1}=3, 2^{-1}=2, 3^{-1}=1$$

\oplus	0	1	2	3
0	0	1	2	3
1	1	2	3	0
2	2	3	0	1
3	3	0	1	2

v) As the table is symmetric w.r.t. to the principal diagonal. Hence commutative law holds.

Ø Question # 3

(i) As $0 \in \mathbb{Q}$, multiplicative inverse of 0 is not in set \mathbb{Q} . Therefore the set of rational number is not a group w.r.t to “ \cdot ”.

(ii) a- Closure property holds in \mathbb{Q} under + because sum of two rational number is also rational.

b- Associative property holds in \mathbb{Q} under addition.

c- $0 \in \mathbb{Q}$ is an identity element.

d- If $a \in \mathbb{Q}$ then additive inverse $-a \in \mathbb{Q}$ such that $a + (-a) = (-a) + a = 0$.

Therefore the set of rational number is group under addition.

(iii) a- Since for $a, b \in \mathbb{Q}^+$, $ab \in \mathbb{Q}^+$ thus closure law holds.

b- For $a, b, c \in \mathbb{Q}$, $a(bc) = (ab)c$ thus associative law holds.

c- Since $1 \in \mathbb{Q}^+$ such that for $a \in \mathbb{Q}^+$, $a \times 1 = 1 \times a = a$. Hence 1 is the identity element.

d- For $a \in \mathbb{Q}^+$, $\frac{1}{a} \in \mathbb{Q}^+$ such that $a \times \frac{1}{a} = \frac{1}{a} \times a = 1$. Thus inverse of a is $\frac{1}{a}$.

Hence \mathbb{Q}^+ is group under addition.

(iv) Since $\mathbb{Z} = \{0, \pm 1, \pm 2, \pm 3, \dots\}$

a- Since sum of integers is an integer therefore for $a, b \in \mathbb{Z}$, $a + b \in \mathbb{Z}$.

b- Since $a + (b + c) = (a + b) + c$ thus associative law holds in \mathbb{Z} .

c- Since $0 \in \mathbb{Z}$ such that for $a \in \mathbb{Z}$, $a + 0 = 0 + a = \mathbb{Z}$. Thus 0 an identity element.

d- For $a \in \mathbb{Z}$, $-a \in \mathbb{Z}$ such that $a + (-a) = (-a) + a = 0$. Thus inverse of a is $-a$.

(v) Since $\mathbb{Z} = \{0, \pm 1, \pm 2, \pm 3, \dots\}$

For any $a \in \mathbb{Z}$ the multiplicative inverse of a is $\frac{1}{a} \notin \mathbb{Z}$. Hence \mathbb{Z} is not a group under multiplication.

Ø Question # 4

As $E + E = E, E + O = O, O + O = E$

Thus the table represents the sums of the elements of set $\{E, O\}$.

\oplus	E	O
E	E	O
O	O	E

The identity element of the set is E because

$$E + E = E + E = E \quad \& \quad E + O = O + E = E .$$

i) From the table each element belong to the set $\{E, O\}$.

Hence closure law is satisfied.

ii) \oplus is associative in $\{E, O\}$

iii) E is the identity element of w.r.t to \oplus

iv) As $O + O = E$ and $E + E = E$, thus inverse of O is O and inverse of E is E .

v) As the table is symmetric about the principle diagonal therefore \oplus is commutative.

Hence $\{E, O\}$ is abelian group under \oplus .

Ø Question # 5

Suppose $G = \{1, w, w^2\}$

\otimes	1	w	w^2
1	1	w	w^2
w	w	w^2	1
w^2	w^2	1	w

i) A table show that all the entries belong to G .

ii) Associative law holds in G w.r.t. multiplication.

e.g. $1 \times (w \times w^2) = 1 \times 1 = 1$

$$(1 \times w) \times w^2 = w \times w^2 = 1$$

iii) Since $1 \times 1 = 1, 1 \times w = w \times 1 = w, 1 \times w^2 = w^2 \times 1 = w^2$

Thus 1 is an identity element in G .

iv) Since $1 \times 1 = 1 \times 1 = 1, w \times w^2 = w^2 \times w = 1, w^2 \times w = w \times w^2 = 1$

therefore inverse of 1 is 1, inverse of w is w^2 , inverse of w^2 is w .

v) As table is symmetric about principle diagonal therefore commutative law holds in G .

Hence G is an abelian group under multiplication.

Ø Question # 6

Given that G is a group under the operation $*$ and $a, b \in G$ such that

$$a * x = b$$

As $a \in G$ and G is group so $a^{-1} \in G$ such that

$$a^{-1} * (a * x) = a^{-1} * b$$

$$\Rightarrow (a^{-1} * a) * x = a^{-1} * b \quad \text{as associative law hold in } G.$$

$$\Rightarrow e * x = a^{-1} * b \quad \text{by inverse law.}$$

$$\Rightarrow x = a^{-1} * b \quad \text{by identity law.}$$

And for

$$x * a = b$$

$$\Rightarrow (x * a) * a^{-1} = b * a^{-1} \quad \text{For } a \in G, a^{-1} \in G$$

$$\Rightarrow x * (a * a^{-1}) = b * a^{-1} \quad \text{as associative law hold in } G.$$

$$\Rightarrow x * e = b * a^{-1} \quad \text{by inverse law.}$$

$$\Rightarrow x = b * a^{-1} \quad \text{by identity law.}$$

Ø Question # 7

Consider $G = \{a + \sqrt{3}b \mid a, b \in \mathbb{Q}\}$

i) Let $a + \sqrt{3}b, c + \sqrt{3}d \in G$, where a, b, c & d are rational.

$$(a + \sqrt{3}b) + (c + \sqrt{3}d) = (a + c) + \sqrt{3}(b + d) = a' + \sqrt{3}b' \in G$$

where $a' = a + c$ and $b' = b + d$ are rational as sum of rational is rational.

Thus closure law holds in G under addition.

ii) For $a + \sqrt{3}b, c + \sqrt{3}d, e + \sqrt{3}f \in G$

$$\begin{aligned} (a + \sqrt{3}b) + ((c + \sqrt{3}d) + (e + \sqrt{3}f)) &= (a + \sqrt{3}b) + ((c + e) + \sqrt{3}(d + f)) \\ &= (a + (c + e)) + \sqrt{3}(b + (d + f)) \\ &= ((a + c) + e) + \sqrt{3}((b + d) + f) \end{aligned}$$

As associative law hold in \mathbb{Q}

$$\begin{aligned} &= ((a + c) + \sqrt{3}(b + d)) + (e + \sqrt{3}f) \\ &= ((a + \sqrt{3}b) + (c + \sqrt{3}d)) + (e + \sqrt{3}f) \end{aligned}$$

Thus associative law hold in G under addition.

iii) $0 + \sqrt{3} \cdot 0 \in G$ as 0 is a rational such that for any $a + \sqrt{3}b \in G$

$$(a + \sqrt{3}b) + (0 + \sqrt{3} \cdot 0) = (a + 0) + \sqrt{3}(b + 0) = a + \sqrt{3}b$$

And $(0 + \sqrt{3} \cdot 0) + (a + \sqrt{3}b) = (0 + a) + \sqrt{3}(0 + b) = a + \sqrt{3}b$

Thus $0 + \sqrt{3} \cdot 0$ is an identity element in G .

iv) For $a + \sqrt{3}b \in G$ where a & b are rational there exist rational $-a$ & $-b$ such that

$$(a + \sqrt{3}b) + ((-a) + \sqrt{3}(-b)) = (a + (-a)) + \sqrt{3}(b + (-b)) = 0 + \sqrt{3} \cdot 0$$

& $((-a) + \sqrt{3}(-b)) + (a + \sqrt{3}b) = ((-a) + a) + \sqrt{3}((-b) + b) = 0 + \sqrt{3} \cdot 0$

Thus inverse of $a + \sqrt{3}b$ is $(-a) + \sqrt{3}(-b)$ exists in G .

v) For $a + \sqrt{3}b, c + \sqrt{3}d \in G$

$$\begin{aligned} (a + \sqrt{3}b) + (c + \sqrt{3}d) &= (a + c) + \sqrt{3}(b + d) \\ &= (c + a) + \sqrt{3}(d + b) \quad \text{As commutative law hold in } \mathbb{Q}. \\ &= (c + d\sqrt{3}) + (a + \sqrt{3}b) \end{aligned}$$

Thus Commutative law holds in G under addition.

And hence G is an abelian group under addition.

Ø Question 8

Let $A, B \in P(S)$ where A & B are subsets of S .

As intersection of two subsets of S is subset of S .

Therefore $A * B = A \cap B \in P(S)$. Thus closure law holds in $P(S)$.

For $A, B, C \in P(S)$

$$A * (B * C) = A \cap (B \cap C) = (A \cap B) \cap C = (A * B) * C$$

Thus associative law holds and $P(S)$.

And hence $(P(S), *)$ is a semi-group.

For $A \in P(S)$ where A is a subset of S we have $S \in P(S)$ such that

$$A \cap S = S \cap A = A.$$

Thus S is an identity element in $P(S)$. And hence $(P(S), *)$ is a monoid.

Ø Question 9

Let x_1 and x_2 be the required elements.

By associative law

$$\begin{aligned} (a * a) * a &= a * (a * a) \\ \Rightarrow c * a &= a * c \\ \Rightarrow x_1 &= b \end{aligned}$$

Now again by associative law

$$\begin{aligned} (a * a) * b &= a * (a * b) \\ \Rightarrow c * b &= a * a \quad \Rightarrow x_2 = c \end{aligned}$$

*	a	b	c
a	c	a	b
b	a	b	c
c	x_1	x_2	a

Question 10

Let G be the all non-singular 2×2 matrices over the real field.

i) Let $A, B \in G$ then $A_{2 \times 2} \times B_{2 \times 2} = C_{2 \times 2} \in G$

Thus closure law holds in G under multiplication.

ii) Associative law in matrices of same order under multiplication holds.

therefore for $A, B, C \in G$

$$A \times (B \times C) = (A \times B) \times C$$

iii) $I_{2 \times 2} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ is a non-singular matrix such that

$$A_{2 \times 2} \times I_{2 \times 2} = I_{2 \times 2} \times A_{2 \times 2} = A_{2 \times 2}$$

Thus $I_{2 \times 2}$ is an identity element in G .

iv) Since inverse of non-singular square matrix exists,

therefore for $A \in G$ there exist $A^{-1} \in G$ such that $AA^{-1} = A^{-1}A = I$.

v) As we know for any two matrices $A, B \in G$, $AB \neq BA$ in general.

Therefore commutative law does not holds in G under multiplication.

Hence the set of all 2×2 non-singular matrices over a real field is a non-abelian group under multiplication.

Made by: **Atiq ur Rehman** (mathcity@gmail.com),
<http://www.mathcity.org>

The End

QUOTATIONS

▣ The tragedy of life is not that it ends so soon, but that we wait so long to begin it.
W. M. Lewis

▣ It is neither good nor bad, but thinking makes it so.
William Shakespeare

▣ Why fear death? It is the most beautiful adventure in life.
Charles Frohman

▣ An unfortunate thing about this world is that the good habits are much easier to give up than the bad ones.
W. Somerset Maugham

▣ The truth is that life is delicious, horrible, charming, frightful, sweet, bitter, and that it is everything.
Anatole France

▣ There is big difference between getting ready to act and starting to act. Many are forever ready to act.
Anonymous